

# Race and Bias in Child Maltreatment Diagnosis and Reporting

Vincent J. Palusci, MD, MS, FAAP,<sup>a</sup> Ann S. Botash, MD, FAAP<sup>b</sup>

Pediatricians have implicit and explicit racial biases that impact the health and well-being of children and their families.<sup>1,2</sup> Similarly, effects of racism on diagnosis and reporting of suspected child abuse and neglect to child protective services (CPS) can have serious consequences. Although we and others are mandated to report suspected child abuse or neglect in all US states and territories, the threshold for reporting requires only “reasonable suspicion” of abuse or neglect.<sup>3</sup> Pediatricians may also report families that they perceive need additional resources.

Research on racial bias and physician reporting is limited.<sup>4</sup> Disproportionality in reporting is widespread on the basis of race, culture, and ethnicity. Asian American and Hispanic children are underrepresented, but overrepresentation of Black children throughout the child protection system has been most widely described.<sup>5</sup> Black children are reported at approximately twice the rate of white children, and the complex relationship of reporting with poverty and race has yet to be fully understood.<sup>4-6</sup> Once reported, cases with Black children are more likely to be accepted for investigation, be confirmed, be brought to court, result in removal of the children from their families for longer periods of time, and take longer

to be closed, possibly related to surveillance bias.<sup>5,7</sup> Multiple points in this process are subject to bias, but the process begins with reporting.<sup>8,9</sup>

The concept of bias has been recognized and discussed within the medical community for at least half a century. Diagnostic errors due to bias can lead to errors in reporting.<sup>10</sup> Although physicians have historically been key players in the development of US reporting systems, they and other medical sources accounted for only 10.5% of the 4.3 million reports to CPS in 2018.<sup>11</sup> Even so, CPS and other agencies rely heavily on medical diagnosis from health care providers, so biases can be magnified.<sup>12,13</sup> In 1985, Hampton and Newberger<sup>14</sup> showed hospitals failed to report almost one-half of cases meeting their study’s definition of abuse. White children and those with higher family income and employment had fewer reports.

Studies have highlighted bias in the medical evaluation of child abuse and neglect. In a classic report, Jenny et al<sup>15</sup> found 31.2% of 173 children with abusive head trauma had been previously seen by physicians. The diagnosis was more likely to be missed in young children from white, “intact” families. Worse, 27.8% were reinjured after the missed diagnosis, and 4 of 5 deaths

<sup>a</sup>Department of Pediatrics, Grossman School of Medicine, New York University, New York, New York; and <sup>b</sup>Department of Pediatrics, State University of New York Upstate Medical University, Syracuse, New York

Drs Palusci and Botash drafted the initial manuscript, reviewed and revised the manuscript, approved the manuscript as submitted, and agree to be accountable for all aspects of the work.

**DOI:** <https://doi.org/10.1542/peds.2020-049625>

Accepted for publication Mar 9, 2021

Address correspondence to Vincent J. Palusci, MD, MS, FAAP, Department of Pediatrics, Grossman School of Medicine, New York University, 462 First Ave, Room GC-65, New York, NY 10016. E-mail: [vincent.palusci@nyulangone.org](mailto:vincent.palusci@nyulangone.org)

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275)

Copyright © 2021 by the American Academy of Pediatrics

**FINANCIAL DISCLOSURE:** The authors have indicated they have no financial relationships relevant to this article to disclose.

**FUNDING:** Supported in part by funding for the New York State Department of Health Child Abuse Medical Provider Program.

**POTENTIAL CONFLICT OF INTEREST:** Dr Palusci is a consultant to the New York City Administration for Children and has acted as a paid consultant to attorneys. Dr Botash is the indirect recipient of grant funding as director of the Child Abuse Medical Provider Program funded by the New York State Department of Health under contract C-505335.

**To cite:** Palusci V J, Botash A S. Race and Bias in Child Maltreatment Diagnosis and Reporting. *Pediatrics*. 2021;148(1):e2020049625

might have been prevented. Lane et al<sup>16</sup> later found that underrepresented minority children seen for fractures were more likely to be evaluated with a skeletal survey and reported, even after controlling for insurance status, independent expert determination, and likelihood of abusive injury. They concluded, "It is possible biases on the part of mandated reporters may contribute to these differences."

Others identified similar trends. Wood et al<sup>17</sup> noted fewer skeletal surveys among white infants evaluated for traumatic brain injury. Hymel et al<sup>18</sup> observed significant race- and ethnicity-based disparities in abusive head trauma evaluation and reporting, concluding, "in the absence of local confounders, these disparities likely represent the impact of local physicians' implicit bias." Reports from medical providers have also been found to be affected by socioeconomic status, ethnicity, and culture.<sup>19,20</sup>

Diagnostic errors in medicine are influenced by knowledge gaps, communication skills, and access to resources, as well as cognitive and implicit biases.<sup>21,22</sup> An Institute of Medicine report<sup>23</sup> describes clinical reasoning as "the cognitive process that is necessary to evaluate and manage a patient's medical problems" and, in addition, describes a dual process theory, one analytical and one nonanalytical. Systems I thinking is described as intuitive, automatic, and based on pattern recognition by using rules of thumb and mental shortcuts. In comparison, systems II thinking uses metacognition and is deliberate and analytical, with hypothetical and/or deductive reasoning and logic.

Skellern<sup>24</sup> describes examples of errors in both types of reasoning applied to child abuse. Systems I errors in diagnosis may be difficult

to overcome because these are unconscious processes.<sup>22</sup> Mandated reporting requires only suspicion, and pediatricians may rely on their "gut" and systems I thinking, potentially relying on implicit biases. Stolper et al<sup>25</sup> report on the use of "gut feelings," the further steps needed to analyze the source of suspicions of abuse, and their need for more training. Gut feelings regarding abuse, potentially triggered by systems I thinking and potentially biased, may initiate the pediatrician's diagnosis and suspicion for reporting. These thoughts and feelings should be followed by analytical reasoning using systems II thinking.<sup>26</sup>

It is important for pediatricians in all practice settings to be able to make appropriate mandated reports for suspected child maltreatment. We offer some suggestions for minimizing the effects of bias in reporting, with the goal of equitably improving the safety of all children.

First, pediatricians need ongoing education with comprehensive information about definitions, examples, and indicators of the major types of childhood maltreatment, racial disproportionality, and systemic and implicit biases. A recent study highlighted gaps in existing training curricula regarding identifying child maltreatment and when and why reports should be made.<sup>27</sup>

Second, standard tools, such as screening protocols, clinical guidelines, and electronic health records that trigger reportable concerns independent of clinician judgment may help to objectively inform decisions.<sup>28,29</sup> However, as our experience with neonatal drug testing protocols teaches us, these tools have to be nonbiased and instituted with adequate training and compliance coupled with attitudinal change to be

effective.<sup>30,31</sup> Paper checklists and published screening tools can also provide support for pediatricians' systems II thinking.<sup>26,32</sup>

Third, we should strive to build on efforts to develop strong multidisciplinary teams that use diverse cultural and racial perspectives.<sup>33</sup> Interprofessional teamwork, recommended in hospitals, must be intentionally used for the purpose of reflecting on bias.<sup>34</sup> There are many hospital child protection teams and community-based child advocacy centers available to help. In addition, physicians practicing in office settings without easy access to teams can also use American Academy of Pediatrics resources to support their work, such as resources from the Council on Child Abuse and Neglect,<sup>35</sup> the Resilience Project,<sup>36</sup> and chapter child abuse committees. Taking a moment to review for bias in each case with others can be a powerful tool.

Fourth, it is important to recognize and reflect on personal knowledge and biases, specifically regarding child abuse and neglect. For example, our interpretation of parenting skills may rely on expectations based on our own experience and cultural and socioeconomic background. Perceptions of family situations, such as when there is use of cultural medicines<sup>37</sup> or teenager parenting,<sup>38</sup> play a role in influencing reporting. In addition, our recognition of bruising and other skin findings may be based on experiences and training in children who have different skin tones.

When considering a mandated report for suspected child abuse or neglect, we recommend considering the following questions to help differentiate systems I from systems II thinking:

- Why do I suspect maltreatment?
- What is the objective evidence?
- If the family does not look like me, share my values, or lives on the “other” side of town, is that affecting my thinking?
- If my gut is telling me to report, why is that?

Similarly, when making a decision not to report, pediatricians should check if they are thinking, “they are such a good family,” “I have known them since they were children,” “my patients would never do that,” or similar emotional reasoning.

Although self-reflection and mitigation of interpersonal bias are important, other factors should be considered, such as the many systemic and structural forces at play. Child abuse reporting, diagnosis, management, and treatment necessarily involve multiple community agencies whose work must also be free from bias. The American Academy of Pediatrics<sup>1</sup> has additional suggestions that address some of these forces, including the following: (1) creating a culturally safe clinical environment, (2) training staff in culturally competent care, (3) using strategies that counter or replace negative racial messages and experiences with positive ones, and others.

We all strive for culturally competent care with awareness of bias. Implicit and explicit biases are drivers of disproportionate reporting and investigation. Racial bias can be checked through personal reflection and training, input from a diverse multidisciplinary team, assistance from an objective screening process, and case review by using deductive reasoning methods, such as those provided by child abuse pediatric consultation.

## ACKNOWLEDGMENTS

We thank Amy Caruso Brown, MD, MSc, MSCS, for her review of the article and JoAnne Race, MS, and Trish Booth, MA, for their support of the New York State Child Abuse Medical Provider Program, whose webcasts inspired this article. The New York State Department of Health provides grant funding for the New York State Child Abuse Medical Provider Program, and we appreciate the administrative and fiscal support provided by the staff of the New York State Department of Health and the Research Foundation for the State University of New York on behalf of Upstate Medical University.

## REFERENCES

1. Trent M, Dooley DG, Dougé J; Section on Adolescent Health; Council on Community Pediatrics; Committee on Adolescence. The impact of racism on child and adolescent health. *Pediatrics*. 2019;144(2):e20191765
2. Baetzel AE, Holman A, Dobija N, Reynolds PI, Nafiu OO. Racial disparities in pediatric anesthesia. *Anesthesiol Clin*. 2020;38(2):327–339
3. Children's Bureau, Administration for Children and Families, US Department of Health and Human Services. How to report child abuse and neglect. 2020. Available at: <https://www.childwelfare.gov/topics/responding/reporting/how/>. Accessed May 24, 2021
4. Harris MS. Racial bias as an explanatory factor for racial disproportionality and disparities in child welfare. In: Dettlaff AJ, ed. *Racial Disproportionality and Disparities in the Child Welfare System*. Cham, Switzerland: Springer; 2021:141–158
5. Child Welfare Information Gateway. *Racial Disproportionality and Disparity in Child Welfare*. Washington, DC: US Department of Health and Human Services, Children's Bureau; 2016. Available at: [https://www.childwelfare.gov/pubPDFs/racial\\_disproportionality.pdf](https://www.childwelfare.gov/pubPDFs/racial_disproportionality.pdf). Accessed May 24, 2021
6. Drake B, Jonson-Reid M, Kim H, Chiang C-J, Davalishvili D. Disproportionate need as a factor explaining racial disproportionality in the CW system. In: Dettlaff AJ, ed. *Racial Disproportionality and Disparities in the Child Welfare System*. Cham, Switzerland: Springer; 2021:159–176
7. Roberts DE. Child protection as surveillance of African American families. *J Soc Welf Fam Law*. 2014;36(4):426–437
8. Boatswain-Kyte A, Esposito T, Trocmé N. A longitudinal jurisdictional study of Black children reported to child protection services in Quebec, Canada. *Child Youth Serv Rev*. 2020;116:105219
9. Kokaliari ED, Roy AW, Taylor J. African American perspectives on racial disparities in child removals. *Child Abuse Negl*. 2019;90:139–148
10. O'Sullivan ED, Schofield SJ. Cognitive bias in clinical medicine. *J R Coll Physicians Edinb*. 2018;48(3):225–232
11. US Department of Health and Human Services, Administration for Children and Families, Administration on Children, Youth and Families, Children's Bureau. Child maltreatment. Available at: <https://www.acf.hhs.gov/cb/research-data-technology/statistics-research/child-maltreatment>. Accessed May 24, 2021
12. Palusci VJ, Hicks RA, Vandervort FE. “You are hereby commanded to appear”: pediatrician subpoena and court appearance in child maltreatment. *Pediatrics*. 2001;107(6):1427–1430
13. Johnson JC. Report from the special advisor on equal justice in the New York State courts. 2020. Available at: <https://www.nycourts.gov/whatsnew/pdf/SpecialAdviserEqualJusticeReport.pdf>. Accessed May 24, 2021
14. Hampton RL, Newberger EH. Child abuse incidence and reporting by hospitals: significance of severity, class, and race. *Am J Public Health*. 1985;75(1):56–60
15. Jenny C, Hymel KP, Ritzen A, Reinert SE, Hay TC. Analysis of missed cases of abusive head trauma. *JAMA*. 1999;281(7):621–626
16. Lane WG, Rubin DM, Monteith R, Christian CW. Racial differences in the evaluation of pediatric fractures for

- physical abuse. *JAMA*. 2002;288(13):1603–1609
17. Wood JN, Hall M, Schilling S, Keren R, Mitra N, Rubin DM. Disparities in the evaluation and diagnosis of abuse among infants with traumatic brain injury. *Pediatrics*. 2010;126(3):408–414
  18. Hymel KP, Laskey AL, Crowell KR, et al; Pediatric Brain Injury Research Network (PediBIRN) Investigators. Racial and ethnic disparities and bias in the evaluation and reporting of abusive head trauma. *J Pediatr*. 2018;198:137–143.e1
  19. Laskey AL, Stump TE, Perkins SM, Zimet GD, Sherman SJ, Downs SM. Influence of race and socioeconomic status on the diagnosis of child abuse: a randomized study. *J Pediatr*. 2012;160(6):1003–1008.e1
  20. Palusci VJ, Rosman I, Shami MR, Skelnar D. Hospital experience using cultural interpreters with the orthodox Jewish community. *Int J Child Health Hum Dev*. 2020;13(4):415–24
  21. Singh H, Schiff GD, Graber ML, Onakpoya I, Thompson MJ. The global burden of diagnostic errors in primary care. *BMJ Qual Saf*. 2017;26(6):484–494
  22. Zwaan L, Monteiro S, Sherbino J, Ilgen J, Howey B, Norman G. Is bias in the eye of the beholder? A vignette study to assess recognition of cognitive biases in clinical case workups. *BMJ Qual Saf*. 2017;26(2):104–110
  23. National Academies of Sciences, Engineering, and Medicine. *Improving Diagnosis in Health Care*. Washington, DC: The National Academies Press; 2015
  24. Skellern C. Thinking fast and slow in the evaluation of injury plausibility in child protection. *J Paediatr Child Health*. 2020;56(9):1330–1334
  25. Stolper E, van de Wiel M, Kooijman S, Feron F. How child health care physicians struggle from gut feelings to managing suspicions of child abuse. *Acta Paediatr*. 2021;110(6):1847–1854
  26. Janson S. Can we trust our gut feeling when we suspect child abuse? *Acta Paediatr*. 2021;110(6):1713–1714
  27. Baker AJL, LeBlanc S, Adebayo T, Mathews B. Training for mandated reporters of child abuse and neglect: content analysis of state-sponsored curricula. *Child Abuse Negl*. 2021;113:104932
  28. Doubeni CA, Simon M, Krist AH. Addressing systemic racism through clinical preventive service recommendations from the US Preventive Services Task Force. *JAMA*. 2021;325(7):627–628
  29. Rumball-Smith J, Fromkin J, Rosenthal B, et al. Implementation of routine electronic health record-based child abuse screening in General Emergency Departments. *Child Abuse Negl*. 2018;85(11):58–67
  30. Kerker BD, Horwitz SM, Leventhal JM. Patients' characteristics and providers' attitudes: predictors of screening pregnant women for illicit substance use. *Child Abuse Negl*. 2004;28(2):209–223
  31. Ellsworth MA, Stevens TP, D'Angio CT. Infant race affects application of clinical guidelines when screening for drugs of abuse in newborns. *Pediatrics*. 2010;125(6). Available at: [www.pediatrics.org/cgi/content/full/125/6/e1379](http://www.pediatrics.org/cgi/content/full/125/6/e1379)
  32. Greiner MV, Palusci VJ, Keeshin BR, Kearns SC, Sinal SH. A preliminary screening instrument for early detection of medical child abuse. *Hosp Pediatr*. 2013;3(1):39–44
  33. Herbert JL, Bromfield L. Evidence for the efficacy of the Child Advocacy Center model: a systematic review. *Trauma Violence Abuse*. 2016;17(3):341–357
  34. Baird J, Ashland M, Rosenbluth G. Inter-professional teams: current trends and future directions. *Pediatr Clin North Am*. 2019;66(4):739–750
  35. American Academy of Pediatrics. Child abuse and neglect. Available at: <https://www.aap.org/en-us/advocacy-and-policy/aap-health-initiatives/Child-Abuse-and-Neglect/Pages/Child-Abuse-and-Neglect.aspx>. Accessed March 4, 2021
  36. American Academy of Pediatrics. The resilience project: child abuse and neglect. Available at: <https://www.aap.org/en-us/advocacy-and-policy/aap-health-initiatives/resilience/Pages/Child-Abuse-and-Neglect.aspx>. Accessed March 4, 2021
  37. Look KM, Look RM. Skin scraping, cupping, and moxibustion that may mimic physical abuse. *J Forensic Sci*. 1997;42(1):103–105
  38. Weed K, Nicholson JS. Differential social evaluation of pregnant teens, teen mothers and teen fathers by university students. *Int J Adolesc Youth*. 2015;20(1):1–16